

**CS513, Spring 11**  
**Prof. Ron**

**Assignment #6**

**Due May 5, 2011**

(1) Write a routine that, given a matrix  $A_{m \times m}$ , and a number  $n < m$ , computes the matrix  $Q_n$  and the matrix  $\tilde{S}_n$  that appear in the Arnoldi Method (which is the algorithm that was presented in class on Tuesday, April 19). Note that  $\tilde{S}_n$  is  $(n + 1) \times n$ . Make it square by deleting its last row. Denote this new matrix by  $S'_n$ .

Then choose a random matrix  $A$  of order  $100 \times 100$ . Using `matlab` find the spectrum of this matrix, and graph the absolute values of the eigenvalues.

Compare now  $\sigma(A)$  to  $\sigma(S'_n)$ , for different (small) values of  $n$ . Draw conclusions concerning (i) how fast  $\sigma(S'_n)$  converges to  $\sigma(A)$ , and (ii) the "type" of eigenvalues in  $\sigma(A)$  which are actually approximated by  $\sigma(S'_n)$  (for a small  $n$ ).

Turn in your code, the output of your experiment, and your observations.

(2) Let  $I_1, I_2, I_3$  be three intervals of length  $\varepsilon$  centered around 1, 5 and 10 respectively. Choose randomly 100 eigenvalues from the union of the three intervals, and create a matrix  $A$  whose spectrum are these 100 numbers (first, create a diagonal matrix  $D$  with the above eigenvalues, and then define  $A = PDP^{-1}$ , for some random, well-conditioned,  $P$ .) Let  $b \in \mathbb{R}^{100}$  be a randomly chosen vector. Use your routine in Q. 1 in order to write a routine that implements the GMRES method for solving the system  $Ax = b$ :

(i) Use first `matlab` and find  $x$  directly

(ii) Apply GMRES for  $n = 5, 10, 20, 90$ . For each case, check accuracy.

(iii) Turn in your code, the output, and your observations. Try to include in your observations the speed of convergence, and to compare, if possible, to the theory studied in class.

Note: there is a parameter  $\varepsilon$  here. You can play with  $\varepsilon$  and see how its choice affects the speed of convergence.

(3) In this question, you investigate the power of the SVD.

(i) Create a random  $30 \times 10$  matrix  $B$ . Then, multiply your matrix  $B$  by a matrix  $C_{10 \times 10}$ , such that the entries in the first two rows of  $C$  are, say,  $k$  times larger than the entries in the other rows. Denote  $A = BC$ .

(ii) Use the SVD in order to find the two dominant singular vectors of  $A$  (i.e., the first two columns of  $U$  in  $A = U\Sigma V'$ ). Try to draw connections between  $A$  and those singular vectors. Note that your experiment involves the parameter  $k$ , so make sure you run a few experiments with different values of  $k$ .