COMPUTING THE RAMANUJAN TAU FUNCTION

DENIS XAVIER CHARLES

ABSTRACT. We show that the Ramanujan Tau function $\tau(n)$ can be computed by a randomized algorithm in time $O(n^{\frac{1}{2}+\epsilon})$ for every $\epsilon > 0$. The same method also yields a deterministic algorithm that runs in time $O(n^{\frac{3}{4}+\epsilon})$ for every $\epsilon > 0$ to compute $\tau(n)$. Previous algorithms to compute $\tau(n)$ require $\Omega(n)$ time.

1. INTRODUCTION

Let $\tau(n)$ be the coefficient of q^n in the formal expansion $q\prod_{1\leq n}(1-q^n)^{24} = \sum_{1\leq n}\tau(n)q^n$. The following properties of the τ-function are well known:

1. If $n, m \in \mathbb{Z}_{>0}$ such that $\gcd(n,m) = 1$ then $\tau(nm) = \tau(n)\tau(m)$.
2. If $r \geq 1$ and p is a prime then $\tau(p^{r+1}) = \tau(p)\tau(p^r) - p^{11}\tau(p^{r-1})$.

Thus $\tau(n)$ is completely determined by $\tau(p)$ for primes $p|n$. Here is a table of $\tau(p)$ for small prime numbers p.

p	2	3	5	7	11	13
$\tau(p)$	-24	252	4830	-16744	534612	-577738

The importance of the τ-function comes from the fact that it gives the fourier coefficients of a modular form. Namely, the function $\Delta(z) = q\prod_{1\leq n}(1-q^n)^{24}$ where $q = e^{2\pi iz}$ is a cusp form of weight 12 for the full modular group (see [La76]). A famous conjecture of D. H. Lehmer says that $\tau(n)$ is never zero. This conjecture has been verified for all $n \leq 22689242781695999$ [JorKe99]. The function $\tau(n)$ seems to be a hard function to compute. Methods to compute $\tau(n)$ based on recurrence relations that it satisfies or its relations to other arithmetic functions such as $\sigma_k(n)$ require $\Omega(n)$ time steps. Since the number n requires $\log_2 n$ bits these algorithms require exponential time in the length of the input. In this article we show that $\tau(n)$ can be computed in time $O(n^{\frac{1}{2}+\epsilon})$ by a randomized algorithm for every $\epsilon > 0$. Though this algorithm is still an exponential time algorithm it is significantly faster than the other methods. Moreover, algorithms based on recurrences compute values of $\tau(m)$ for $m < n$ when computing $\tau(n)$. Our algorithm has the feature that it does not compute any of the previous values of the τ-function. On the other hand, this algorithm is not well suited to building a table of $\tau(m)$ for all $m < n$ since the table can be built in roughly $O(n)$ time by the other methods, whereas this method would require $O(n^{\frac{3}{2}+\epsilon})$ time. Our algorithm is more suited to computing "spot" values of $\tau(n)$. In the next section we will give the details of the algorithm and prove its running time.

2. THE ALGORITHM

Since we can compute $\tau(n)$ in $O(\log^3 n)$ time provided we know the factorization of the integer n *and* the values of $\tau(p)$ for primes $p|n$, we will concentrate on computing $\tau(p)$ for primes p. There are deterministic algorithms that can factor n in $O(n^{\frac{1}{4}+\epsilon})$ time ([Co93]). We use such an algorithm to find the primes $p|n$. The main idea of the algorithm is to make use of the Selberg Trace formula to compute $\tau(p)$.

Date: March 2003.
Research supported in part by NSF grant CCR-9988202.

Theorem 2.1. [Sel56] *Let $k \geq 4$ be an even integer and let m be an integer > 0. Then the trace of the Hecke operator $T(m)$ on the space of cusp forms $S_k(\Gamma)$ is given by*

$$\text{Tr } T(m) = -\frac{1}{2}\sum_{-\infty<t<\infty} P_k(t,m)H(4m-t^2) - \frac{1}{2}\sum_{dd'=m}\min\{d,d'\}^{k-1}.$$

In the above sum $H(D)$ refers to the Hurwitz class number of D, and $P_k(t,N) = \frac{\rho^{k-1}-\overline{\rho}^{k-1}}{\rho-\overline{\rho}}$ where ρ is a complex number satisfying $\rho+\overline{\rho} = t$ and $\rho\overline{\rho} = N$.

Note that the sum is actually finite since $H(D) = 0$ if $D < 0$ and so if $t > 2\sqrt{m}$, $H(4m-t^2) = 0$.

In our case $\Delta \in S_{12}(\Gamma)$ and it is a one dimensional vector space. The Hecke operators are a family of linear operators $T(n) : S_k(\Gamma) \to S_k(\Gamma)$ for $n \geq 1$ an integer. Since $\dim S_{12}(\Gamma) = 1$, Δ is a simultaneous eigenform for every $T(n)$. It is known (see [La76]) that $T(n)\Delta(z) = \tau(n)\Delta(z)$ where $\Delta(z) \in S_{12}(\Gamma)$ is the function defined earlier. Thus the eigenvalue of the n-th Hecke operator is $\tau(n)$. Since $\dim S_{12}(\Gamma) = 1$, we have $\text{Tr } T(n) = \tau(n)$ and specializing Theorem 2.1 to our case we get the following result:

Theorem 2.2. *Let p be a prime. Then*

$$\tau(p) = -\sum_{0<t\leq\sqrt{4p}} P(t,p)H(4p-t^2) \; + \frac{1}{2}p^5H(4p) \; - 1$$

where

$$P(t,p) = t^{10} - 9t^8p + 28t^6p^2 - 35t^4p^3 + 15t^2p^4 - p^5$$

and $H(D)$ is the Hurwitz class number.

We will use the above theorem to compute $\tau(p)$. In fact, we only need to show how the Hurwitz class numbers can be computed, since it is easy to compute the above sum. For this task we need the following lemma (see [Co93] Lemma 5.3.7):

Lemma 2.3. *Let $w(-3) = 3, w(-4) = 2$ and $w(D) = 1$ for $D < -4$, and set $h'(D) = \frac{h(D)}{w(D)}$, where $h(D)$ is defined to be the class number of the field $\mathbb{Q}(\sqrt{D})$ if $D \equiv 0, 1 \mod 4$ otherwise we define $h(D)$ to be zero. Then for $N > 0$ we have*

$$H(N) = \sum_{d^2|N} h'\left(-\frac{N}{d^2}\right).$$

There are randomized sub-exponential time algorithms to compute the class number (see [Co93]).

Theorem 2.4. *The class number $h(D)$ can be computed deterministically in time $|D|^{\frac{1}{4}+\epsilon}$ for every $\epsilon > 0$, or by a randomized algorithm with expected running time $e^{O(\sqrt{\ln|D|\ln\ln|D|})}$.*

Proposition 2.5. *The Hurwitz class number $H(N)$ can be computed by a deterministic algorithm in time $O(N^{\frac{1}{4}+\epsilon})$ or a randomized algorithm with an expected running time $O(N^{\epsilon})$ for every $\epsilon > 0$.*

Proof : By Lemma 2.3 we have

$$H(N) = \sum_{d^2|N} h'\left(-\frac{N}{d^2}\right).$$

The function $h'(D)$ is essentially just the class number of $\mathbb{Q}(\sqrt{D})$ and so can be computed in time $O(|D|^{\epsilon})$ if we use the randomized algorithm or in time $O(|D|^{\frac{1}{4}+\epsilon})$ if we use the deterministic algorithm. The number of terms in the sum is at most the number of divisors of N. It is known (see [Ten95]) that the number of divisors $d(N) \ll_{\epsilon} N^{\epsilon}$ for every $\epsilon > 0$. Thus the sum can be evaluated by computing each of the terms in the stated time bound. □

Thus putting all these results together we get the following:

Theorem 2.6. *There is a randomized algorithm to compute* $\tau(p)$ *with expected running time* $O(p^{\frac{1}{2}+\epsilon})$ *for every* $\epsilon > 0$.

Theorem 2.7. *There is a deterministic algorithm to compute* $\tau(p)$ *in time* $O(p^{\frac{3}{4}+\epsilon})$ *for every* $\epsilon > 0$.

Acknowledgements: The author would like to thank professor Ken Ono for suggesting the use of the trace formula, and professor Eric Bach for very useful discussions.

References

[Co93] Cohen, Henri; *Computational Algebraic Number Theory*, Graduate Texts in Math. Vol. 138, Springer-Verlag, 1993.

[JorKe99] Jordan, B.; Kelly, B. III; *The vanishing of the Ramanujan Tau function*, Preprint, 1999.

[La76] Lang, Serge; *Introduction to Modular Forms*, Grundlehren der mathematischen Wissenschaften, Vol. 222, Springer-Verlag, 1976.

[Sel56] Selberg, Alte; *Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series*, J. Indian Math. Soc., **20**, 47-87, 1956.

[Ten95] Tenenbaum, Gérald; *Introduction to analytic and probabilistic number theory*, Cambridge Studies in Advanced Mathematics, **46**, Cambridge University Press, 1995.

Department of Computer Science, University of Wisconsin-Madison, Madison WI - 53706.
E-mail address: cdx@cs.wisc.edu