

# CS412 Spring Semester 2011

Midterm #2

Thursday 28 April 2010

Time: 75 mins

Name	
University ID	

Part #1	
Part #2	
Part #3	
Part #4	
Part #5	
TOTAL	

1. [30% = 6 questions  $\times$  5% each] MULTIPLE CHOICE SECTION. Circle or underline the correct answer (or answers). You do not need to provide a justification for your answer(s).
- (1) If the  $n \times n$  matrix  $\mathbf{A}$  is poorly conditioned (i.e. it has a very large condition number), then ...
 

**(Circle or underline the ONE most correct answer)**

    - (a) Solving  $\mathbf{Ax} = \mathbf{b}$  accurately would be difficult with LU decomposition or Gauss elimination, but iterative methods (Jacobi, Gauss-Seidel) would not have a problem.
    - (b) Solving  $\mathbf{Ax} = \mathbf{b}$  accurately with iterative methods (Jacobi, Gauss-Seidel) would be difficult, but LU decomposition with pivoting would not have a problem.
    - (c) Solving  $\mathbf{Ax} = \mathbf{b}$  accurately will be challenging regardless of the method we use.
  - (2) Consider the rectangular  $m \times n$  matrix  $\mathbf{A}$  (with  $m > n$ ) and the vector  $\mathbf{b} \in \mathbf{R}^m$ . If  $\mathbf{x}$  is the *least squares solution* to  $\mathbf{Ax} \approx \mathbf{b}$ , can we say that  $\mathbf{x}$  is an actual solution to  $\mathbf{Ax} = \mathbf{b}$ ?
 

**(Circle or underline the ONE most correct answer)**

    - (a) Yes, in fact  $\mathbf{Ax} = \mathbf{b}$  has many solutions and the “least squares solution” is the one with the smallest 2-norm of the residual vector  $\|\mathbf{r}\|_2$ .
    - (b) No, the system  $\mathbf{Ax} = \mathbf{b}$  will generally not have a solution. What we call the “least squares solution” is the vector  $\mathbf{x}$  with the smallest 2-norm of the error vector  $\|\mathbf{x} - \mathbf{x}_{\text{exact}}\|_2$ .
    - (c) No, the system  $\mathbf{Ax} = \mathbf{b}$  will generally not have a solution. What we call the “least squares solution” is the vector  $\mathbf{x}$  with the smallest 2-norm of the residual vector  $\|\mathbf{b} - \mathbf{Ax}\|_2$ .
  - (3) Which of the following are good reasons for using an iterative method (e.g. Jacobi or Gauss-Seidel) instead of a direct method (e.g. Gauss Elimination or LU factorization) to solve the  $n \times n$  system  $\mathbf{Ax} = \mathbf{b}$ ?
 

**(Circle or underline ALL correct answers)**

    - (a) When an iterative method is convergent and the matrix  $\mathbf{A}$  is relatively sparse, the computational cost of finding a good approximation of the simulation using an iterative approach could be significantly lower than using a direct method.
    - (b) Iterative methods work very well with poorly conditioned matrices, while direct methods face problems in this case.
    - (c) Iterative methods do not require pivoting when  $\mathbf{A}$  is diagonally dominant or symmetric positive definite, while a direct method would require pivoting in this case.

- (4) Imagine that we perform an evaluation of a certain composite integration rule, partitioning the integration interval  $[a, b]$  into equal size subintervals, with length  $h$ . We observe that by doubling the number of data points, the error in the approximation of the integral is reduced by a factor of eight. Which of the following are true?

**(Circle or underline ALL correct answers)**

- (a) The integration rule is third order accurate.
  - (b) The global integration error scales proportionately to  $h^4$ .
  - (c) The local integration error scales proportionately to  $h^4$ .
- (5) When we try to solve an Initial Value Problem  $y' = f(t, y)$ ,  $y(t_0) = y_0$ , why is it desirable for the differential equation to have stable solutions?

**(Circle or underline ALL correct answers)**

- (a) Because in this case numerical methods for approximating the solution will be stable as well.
  - (b) Because in this case it is possible for an propely designed numerical method to match the asymptotic behavior of the exact solution.
  - (c) Because if the solutions were unstable, any errors or inaccuracies incurred at any part of the solution process could be amplified without bound as  $t \rightarrow \infty$ .
- (6) Which of the following statements about norms are true?

**(Circle or underline ALL correct answers)**

- (a)  $\|\mathbf{x}\|_\infty \geq \|\mathbf{x}\|_1$  for all  $\mathbf{x} \in \mathbf{R}^n$  ( $n \geq 2$ ).
- (b)  $\|\mathbf{Ax}\| = \|\mathbf{A}\| \|\mathbf{x}\|$  for any matrix  $\mathbf{A} \in \mathbf{R}^{n \times n}$  and vector  $\mathbf{x} \in \mathbf{R}^n$ .
- (c)  $\|\mathbf{A}^2\| \leq \|\mathbf{A}\|^2$  for any matrix  $\mathbf{A} \in \mathbf{R}^{n \times n}$ .

2. [ $20\% = 4$  questions  $\times 5\%$  each] SHORT ANSWER SECTION. Answer each of the following questions in no more than 1-2 sentences.
- (a) Why do we often prefer to use the composite Simpson's rule, instead of the composite Trapezoidal rule to approximate a definite integral?
  - (b) Write down three ordinary differential equations, one with asymptotically stable solutions, one with stable (but not asymptotically so) solutions, and one with unstable solutions.
  - (c) When solving Initial Value Problems, why does an iteration of an implicit method often require more computational effort, than an iteration of an explicit method?
  - (d) List one of the conditions that would guarantee convergence of the Jacobi method for solving a linear system  $\mathbf{Ax} = \mathbf{b}$ .

3. [16%] Determine the order of accuracy for the following numerical integration rule:

$$\int_a^b f(x)dx \approx \frac{b-a}{2} \left[ f\left(\frac{2a+b}{3}\right) + f\left(\frac{a+2b}{3}\right) \right]$$

4. [14%] Consider the 5 points:

$$\begin{aligned}(x_1, y_1) &= (-3, -1) \\(x_2, y_2) &= (-2, 1) \\(x_3, y_3) &= (0, 2) \\(x_4, y_4) &= (1, 3) \\(x_5, y_5) &= (3, 2)\end{aligned}$$

- (a) We want to determine a straight line  $y = c_1x + c_0$  that approximates these points as closely as possible, in the least squares sense. Write a least squares system  $\mathbf{Ax} \approx \mathbf{b}$  which can be used to determine the coefficients  $c_1$  and  $c_0$ .
- (b) Solve this least squares system, using the method of normal equations.

5. [20%] Consider the following family of methods for solving Initial Value Problems of the form  $y' = f(t, y)$ ,  $y(t_0) = y_0$ :

$$y_{k+1} = y_k + \Delta t [(1-w)f(t_k, y_k) + wf(t_{k+1}, y_{k+1})] \quad (1)$$

In equation (1) the constant  $w$  can take any value in the interval  $[0, 1]$ ; different values produce different methods. We can see, for example, that  $w = 0$  corresponds to Forward Euler,  $w = 0.5$  is Trapezoidal Rule and  $w = 1$  produces Backward Euler.

- (a) Show that for  $0.5 \leq w \leq 1$ , the method of equation (1) is unconditionally stable on the model equation  $y' = \lambda y$ ,  $\lambda < 0$ .
- (b) For  $0 \leq w < 0.5$ , determine the stability condition for the method of equation (1) when applied to the model equation  $y' = \lambda y$ ,  $\lambda < 0$ .

**Hint:** Remember that stability of a method on this model equation is equivalent to showing that  $y_k \rightarrow 0$  as  $k \rightarrow \infty$ .