

CS412 Spring Semester 2013

Practice Midterm #3

1. MULTIPLE CHOICE SECTION. Circle or underline the correct answer (or answers). You do not need to provide a justification for your answer(s).
 - (1) Numerical integration rules which use just one point per interval are:
(Circle or underline the ONE most correct answer)
 - (a) Always first order accurate. Methods that use 2 points are second order accurate and so on.
 - (b) At most first order accurate, but can never have an order of accuracy of 2 or more.
 - (c) Generally first order accurate, but sometimes could also be second order due to fortuitious cancellation.
 - (2) A certain numerical integration rule integrates all cubic polynomials exactly. Which of the following property describes its accuracy?
(Circle or underline the ONE most correct answer)
 - (a) The method is exactly third order accurate.
 - (b) The method is at least fourth order accurate.
 - (c) The global error is proportional to h^3 .
 - (3) Which of the following methods are good choices for solving $\mathbf{Ax} = \mathbf{b}$, where A is symmetric and positive definite?
(Circle or underline ALL correct answers)
 - (a) LU factorization with full pivoting.
 - (b) System of normal equations.
 - (c) Gauss-Seidel method.
 - (d) Jacobi method.
 - (4) If the $n \times n$ matrix \mathbf{A} is poorly conditioned (i.e. it has a very large condition number), then ...
(Circle or underline the ONE most correct answer)
 - (a) Solving $\mathbf{Ax} = \mathbf{b}$ accurately would be difficult with LU decomposition or Gauss elimination, but iterative methods (Jacobi, Gauss-Seidel) would not have a problem.
 - (b) Solving $\mathbf{Ax} = \mathbf{b}$ accurately with iterative methods (Jacobi, Gauss-Seidel) would be difficult, but LU decomposition with pivoting would not have a problem.
 - (c) Solving $\mathbf{Ax} = \mathbf{b}$ accurately will be challenging regardless of the method we use.

- (5) Imagine that we perform an evaluation of a certain composite integration rule, partitioning the integration interval $[a, b]$ into equal size subintervals, with length h . We observe that by doubling the number of data points, the error in the approximation of the integral is reduced by a factor of eight. Which of the following are true?
(Circle or underline ALL correct answers)
- (a) The integration rule is third order accurate.
 - (b) The global integration error scales proportionately to h^4 .
 - (c) The local integration error scales proportionately to h^4 .
- (6) Which of the following are good reasons for using an iterative method (e.g. Jacobi or Gauss-Seidel) instead of a direct method (e.g. Gauss Elimination or LU factorization) to solve the $n \times n$ system $\mathbf{Ax} = \mathbf{b}$?
(Circle or underline ALL correct answers)
- (a) When an iterative method is convergent and the matrix \mathbf{A} is relatively sparse, the computational cost of finding a good approximation of the solution using an iterative approach could be significantly lower than using a direct method.
 - (b) Iterative methods work very well with poorly conditioned matrices, while direct methods face problems in this case.
 - (c) Iterative methods do not require pivoting when \mathbf{A} is diagonally dominant or symmetric positive definite, while a direct method would require pivoting in this case.
- (7) Consider the rectangular $m \times n$ matrix \mathbf{A} (with $m > n$) and the vector $\mathbf{b} \in \mathbf{R}^m$. If \mathbf{x} is the *least squares solution* to $\mathbf{Ax} \approx \mathbf{b}$, can we say that \mathbf{x} is an actual solution to $\mathbf{Ax} = \mathbf{b}$?
(Circle or underline the ONE most correct answer)
- (a) Yes, in fact $\mathbf{Ax} = \mathbf{b}$ has many solutions and the “least squares solution” is the one with the smallest 2-norm of the residual vector $\|\mathbf{r}\|_2$.
 - (b) No, the system $\mathbf{Ax} = \mathbf{b}$ will generally not have a solution. What we call the “least squares solution” is the vector \mathbf{x} with the smallest 2-norm of the error vector $\|\mathbf{x} - \mathbf{x}_{\text{exact}}\|_2$.
 - (c) No, the system $\mathbf{Ax} = \mathbf{b}$ will generally not have a solution. What we call the “least squares solution” is the vector \mathbf{x} with the smallest 2-norm of the residual vector $\|\mathbf{b} - \mathbf{Ax}\|_2$.

2. SHORT ANSWER SECTION. Answer each of the following questions in no more than 1-2 sentences.

- (a) Describe one scenario where you would prefer using the **LU** factorization, instead of an iterative method such as Jacobi or Gauss-Seidel, for solving a linear system $\mathbf{Ax} = \mathbf{b}$.

- (b) What is one benefit of the **QR** factorization when compared to the normal equations, as methods for solving least squares problems?

- (c) Describe one reason why solving a system $\mathbf{Ax} = \mathbf{b}$ could be extremely challenging when \mathbf{A} has a very high condition number.

- (d) List one of the conditions that would guarantee convergence of the Jacobi method for solving a linear system $\mathbf{Ax} = \mathbf{b}$.

- (e) Why is Simpson's rule potentially much more attractive than the trapezoidal rule, when approximating definite integrals?

- (f) Describe one plausible stopping criterion for determining when to stop an iterative solver for $\mathbf{Ax} = \mathbf{b}$, such as Jacobi or Gauss-Seidel.

3. Consider the $n > 3$ data points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$. We want to find a cubic polynomial $p(x) = c_3x^3 + c_2x^2 + c_1x + c_0$ such that the graph of $p(x)$ approximates the given data points as much as possible. Write a Least Squares system $\mathbf{Ax} \approx \mathbf{b}$ which can be used to determine this cubic approximating polynomial. What does this system reduce to, in the case $n = 4$?
4. Show that the coefficient matrix in the system of normal equations (used for solving least squares problems) is always symmetric and positive definite.
5. Consider the $n \times n$ linear system $\mathbf{Ax} = \mathbf{b}$.
- Show that if \mathbf{A} is diagonal, the Jacobi method converges after just one iteration.
 - Show that if \mathbf{A} is lower triangular, the Gauss-Seidel method converges after just one iteration.
6. Consider the Ordinary Partial Differential Equation

$$y'(t) = -\sin^2(t)y^3(t)$$

- Write the formula for the Forward Euler method applied to this differential equation
 - Write the formula for the Backward Euler method applied to this differential equation.
 - Write the iterative formula for Newton's method, used to solve the equation derived in question (b) above.
7. The numerical integration rule known as *Milne's Rule* is defined as:

$$\int_a^b f(x)dx \approx \frac{b-a}{3} \left[2f\left(\frac{3a+b}{4}\right) - f\left(\frac{a+b}{2}\right) + 2f\left(\frac{a+3b}{4}\right) \right]$$

Determine the order of accuracy of this method.

8. Consider a numerical integration rule defined as:

$$\int_a^b f(x)dx \approx w_1f(a) + w_2f\left(\frac{2a+b}{3}\right) + w_3f(b)$$

where w_1, w_2, w_3 are undetermined constants. Find the values of w_1, w_2, w_3 such that this rule becomes 3rd order accurate.