

EQUI-PARTITIONING OF HIGHER-DIMENSIONAL HYPER-RECTANGULAR GRID GRAPHS *

ATHULA GUNAWARDENA [†] AND ROBERT R. MEYER[‡]

Abstract. A d -dimensional grid graph G is the graph on a finite subset in the integer lattice Z^d in which a vertex $x = (x_1, x_2, \dots, x_n)$ is joined to another vertex $y = (y_1, y_2, \dots, y_n)$ if for some i we have $|x_i - y_i| = 1$ and $x_j = y_j$ for all $j \neq i$. G is *hyper-rectangular* if its set of vertices forms $[K_1] \times [K_2] \times \dots \times [K_d]$, where each K_i is a nonnegative integer, $[K_i] = \{0, 1, \dots, K_i - 1\}$. The *surface area* of G is the number of edges between G and its complement in the integer grid Z^d . We consider the Minimum Surface Area problem, $MSA(G, V)$, of partitioning G into subsets of cardinality V so that the total surface area of the subgraphs corresponding to these subsets is a minimum. Although several efficient 2-dimensional heuristics which exploit the geometry of the grid are available in the literature, very little progress has been made in constructing similar algorithms for partitioning higher dimensional ($d > 2$) grid graphs. We present an equi-partitioning algorithm for higher dimensional hyper-rectangles and establish related asymptotic optimality properties. Our algorithm generalizes the two dimensional algorithm due to Martin [*Discrete Appl. Math.*, 82 (1998), pp. 193–207]. It runs in linear time in the number of nodes ($O(n), n = |G|$) when each K_i is $O(n^{1/d})$. Utilizing a result due to Bollabas and Leader [Combinatorica, 11.4 (1991), pp. 299–314], we construct an algorithm to calculate a useful lower bound for the surface area of an equi-partition. Our computational results either achieve this lower bound (i.e., are optimal) or stay within a few percent of the bound.

Key words. higher dimensional grid graph, hyper-rectangle, graph partitioning, knapsack problem

AMS subject classifications. 05A18, 68R01, 68W01

1. Introduction. Grid graph equi-partition arises in the context of minimizing interprocessor communication subject to load balancing in parallel computation for a variety of problem classes including the solution of PDEs using finite difference schemes [11], computer vision [10], and database applications [7]. Similar to the general graph partitioning problem [6], arbitrary grid graph partitioning problem is NP-complete [12], and at present one has to depend on effective heuristic algorithms.

In this paper, we focus on partitioning graph analogs of hyper-rectangles and hereafter call our problem the Minimum Surface Area problem for rectilinear grid graphs, denoted by $MSA(G, V)$, in which the d -dimensional grid graph $G = [K_1] \times [K_2] \dots \times [K_d]$ is partitioned into subgraphs of cardinality V so that the total surface area of the subgraphs is a minimum. We assume that V divides $K_1 K_2 \dots K_d$. Figure 1.1 shows a 25-component equi-partition for the $MSA([5]^3, 5)$ problem generated by our equi-partitioning algorithm presented in Section 4. Since the surface area of this partition does not equal the lower bound derived in Section 3, its optimality is not guaranteed but its relative optimality gap $[(\text{area} - \text{lower bound}) / \text{lower bound}] (= 2 / 500)$ is 0.04%, and only one component allows any possibility for surface area improvement.

Several effective algorithms for partitioning a 2-dimensional grid graph are available in the literature [2, 4, 5, 8, 14]. Our algorithm generalizes the fast two dimensional algorithm due to Martin [8]. With this generalization, we reduce the problem to several knapsack problems and use dynamic programming to solve them.

*

[†]Department of Computer Science, University of Wisconsin-Madison, Wisconsin 53706 (agunawardena@wisc.edu), and Department of Mathematical and Computer Sciences, University of Wisconsin-Whitewater, Whitewater, Wisconsin 53190 (gunawara@uw.edu).

[‡]Department of Computer Science, University of Wisconsin-Madison, Wisconsin 53706 (rrm@cs.wisc.edu). The research of this author was supported in part by NSF grant DMI-0100220 .

V	V	T	R	P
V	V	T	R	P
L	L	J	H	F
L	L	J	H	F
D	D	B	B	A

V	V	T	R	P
W	W	T	R	P
L	M	J	H	F
M	M	J	H	F
D	D	B	B	A

X	X	U	S	Q
W	W	T	R	P
N	N	K	I	G
M	M	J	H	F
D	E	B	C	A

X	X	U	S	Q
X	Y	U	S	Q
N	N	K	I	G
N	O	K	I	G
E	E	C	C	A

Y	Y	U	S	Q
Y	Y	U	S	Q
O	O	K	I	G
O	O	K	I	G
E	E	C	C	A

FIG. 1.1. An equi-partition for $MSA([5]^3, 5)$ with bound gap 0.04%. Only component A does not have minimum surface area.

Those grid graphs that meet the minimum surface area for a given number of vertices draw special interest from research in Combinatorics (i.e., study of Polyominoes) [1], and in Physics (i.e., study of the metastable behavior of the stochastic ising model) [9]. In 1991, Bollabas and Leader [3] derived a class of grid graphs with minimum surface area. In Section 3, utilizing their result, we provide a formula for the minimum surface area for grid graphs with a given cardinality and use this result to find a lower bound for the surface area of a given equi-partition. We present our equi-partitioning algorithm in Section 4 and give computational results in Section 5 for some 3 and 4-dimensional equi-partitions. We also compare our results with the lower bound given in Section 3.

2. Preliminaries. We follow the notation and definitions given in [3]. For the readers' convenience, we restate them here. A d -dimensional grid graph is the graph on a finite subset in the integer lattice Z^d in which $x = (x_1, x_2, \dots, x_n)$ is joined to $y = (y_1, y_2, \dots, y_n)$ if for some i we have $|x_i - y_i| = 1$ and $x_j = y_j$ for all $j \neq i$. Since grid graphs are determined by their nodes, in much of the discussion below we identify the graph with its nodes. Let $P([n])$ be the power set of $[n] = \{0, 1, \dots, n-1\}$. The *binary order* on $P([n])$ is a total order on $P([n])$'s 2^n elements such that for some $S, T \in P$, $S < T$ if and only if the greatest element of $[n]$ which is in one of S and T but not the other is actually in T (i.e., $\max(S \Delta T) \in T$). The *cube order* on $[k]^n$ is a total order such that for some $x = (x_1, x_2, \dots, x_n)$ and $y = (y_1, y_2, \dots, y_n)$, $x < y$ if and only if for some $s \in [k]$, $\{i : x_i = s\} < \{i : y_i = s\}$ in the binary order on $P([n])$ with $\{i | x_i = t\} = \{i | y_i = t\}$ for all $t > s$. For $S \subseteq [k]^n$, the cube order on S is the restriction to S of the cube order on $[k]^n$. We write $x + [k]$ to represent $\{x, x+1, \dots, x+k-1\}$.

Let G be a grid graph and E be the set of edges in Z^n . The *edge boundary* of G , denoted by $\sigma(G)$, is the set $\{(x, y) \in E | x \in G, y \notin G\}$. The *surface area* of G , denoted by $\mathcal{A}(G)$, is the cardinality of $\sigma(G)$ ($|\sigma(G)|$) (In three dimensional space this matches the "exposed" surface area of the set of unit hypercubes with centers at the nodes of G). The following proposition states that the cube order may be used to create graphs with minimum surface area. This is a direct corollary to Theorem 15 in [3]. Other versions of proofs of this result can be found in [9], [1] ($d = 3$), and [13].

PROPOSITION 2.1. *Let G be a d -dimensional grid graph such that $G \subseteq [k]^d$ for some large enough positive integer k , and J be the set of first $|G|$ elements in the cube order on $[k]^d$. Then $\mathcal{A}(G) \geq \mathcal{A}(J)$.*

Let d be a positive integer and n_d be a nonnegative integer. Let k_d be the unique integer such that $k_d^d \leq n_d < (k_d + 1)^d$, and m_d be the unique integer in $[d]$ such that $(k_d + 1)^{m_d} k_d^{d-m_d} \leq n_d < (k_d + 1)^{m_d + 1} (k_d)^{d-m_d-1}$. Observe that $k_1 = n_1$ and $m_1 = 0$.

For an integer n_d , we define the following special grid graph, called the *quasi-cube* recursively, and we show in Section 3 that quasi-cubes are generated by the cube order.

DEFINITION 2.2 (Quasi-Cube). *The d -dimensional quasi-cube corresponding to n_d is denoted by $C_d(n_d)$. For $d = 1$, $C_1(n_1) = [n_1]$. For $d > 1$, $C_d(n_d)$ is the set*

$$[k_d + 1]^{m_d} \times [k_d]^{d-m_d} \cup P(C_{d-1}(n_{d-1})),$$

where k_d and m_d are unique nonnegative integers as mentioned above, $n_{d-1} = n_d - (k_d + 1)^{m_d} k_d^{d-m_d}$, and $P(C_{d-1}(n_{d-1}))$ is the set of points, $(x_1, x_2, \dots, x_{m_d}, k_d, x_{m_d+1}, \dots, x_{d-1})$, such that $(x_1, x_2, \dots, x_{d-1}) \in C_{d-1}(n_{d-1})$.

Since k_i and m_i ($d \geq i \geq 1$) are unique integers, the quasi-cube $C_d(n_d)$ is uniquely defined.

3. A Lower Bound for the Total Surface Area of an Equi-Partition.

We use the quasi-cube to obtain our results in this section. The following lemmas describe some of the properties of the quasi-cube.

LEMMA 3.1. *The quasi-cube has the properties:*

$$(a) |C_d(n_d)| = n_d = \sum_{r=1}^d (k_r + 1)^{m_r} k_r^{r-m_r}.$$

$$(b) \mathcal{A}(C_d(n_d)) = 2 \sum_{r=1}^d [(r - m_r)(k_r + 1)^{m_r} k_r^{r-m_r-1} + m_r(k_r + 1)^{m_r-1} k_r^{r-m_r}].$$

Proof. By Definition 2.2, $C_d(n_d) = S_d \cup P(C_{d-1}(n_{d-1}))$ where $S_d = [k_d + 1]^{m_d} \times [k_d]^{d-m_d}$. It is clear that S_d and $P(C_{d-1}(n_{d-1}))$ are disjoint and $|P(C_{d-1}(n_{d-1}))| = |C_{d-1}(n_{d-1})|$. So we have $|C_d(n_d)| = |S_d| + |C_{d-1}(n_{d-1})|$, and get (a) by applying induction on d .

To prove (b), we consider the surface area of $C_d(n_d)$. We have $\mathcal{A}(C_d(n_d)) = \mathcal{A}(S_d) + \mathcal{A}(P(C_{d-1}(n_{d-1}))) - 2|\sigma(S_d) \cap \sigma(P(C_{d-1}(n_{d-1})))|$. Since $k_{d-1} \leq k_d$, we get $\sigma(P(C_{d-1}(n_{d-1}))) \subseteq \sigma(S_d)$. Using $\mathcal{A}(S_d) = 2(d - m_d)(k_d + 1)^{m_d} k_d^{d-m_d-1} + 2m_d(k_d + 1)^{m_d-1} k_d^{d-m_d}$, $\mathcal{A}(P(C_{d-1}(n_{d-1}))) = \mathcal{A}(C_{d-1}(n_{d-1})) + 2|C_{d-1}(n_{d-1})|$, and $|\sigma(S_d) \cap \sigma(P(C_{d-1}(n_{d-1})))| = |C_{d-1}(n_{d-1})|$, we get the recurrence relation $\mathcal{A}(C_d(n_d)) = \mathcal{A}(C_{d-1}(n_{d-1})) + 2(d - m_d)(k_d + 1)^{m_d} k_d^{d-m_d-1} + 2m_d(k_d + 1)^{m_d-1} k_d^{d-m_d}$, and the initial condition $\mathcal{A}(C_1(n_1)) = 2$ if $n_1 > 0$ and, 0 if $n_1 = 0$. It is straightforward to solve this recurrence relation and get our result in (b). \square

LEMMA 3.2. *The quasi-cube, $C_d(n_d)$, has the minimum surface area among the d -dimensional grid graphs with n_d vertices.*

Proof. First we prove that the quasi-cube $C_d(n_d)$ forms an initial segment of the cube order in dimension d . We prove this by induction on d . Clearly $C_1(n_1) = [n_1]$ is an initial segment of the cube order. Assume $C_{d-1}(n_{d-1})$ is an initial segment of the cube order. By the definition, $C_d(n_d) = S_d \cup P(C_{d-1}(n_{d-1}))$ where $S_d = [k_d]^{m_d} \times [k_d + 1]^{d-m_d}$. Let $\bar{x} = (x_1, x_2, \dots, x_d)$, where $x_i = k_d - 1$ if $i \leq m_d$ and $x_i = k_d$ if $i > m_d$. Clearly $\bar{x} \in S_d$ and any element y in the cube order such that $y < \bar{x}$ is in S_d . Since there are $|S_d|$ elements which are less than or equal to \bar{x} in cube order, S_d is an initial segment of the cube order. We may assume that $C_{d-1}(n_{d-1}) \neq \phi$. Otherwise $C_d(n_d) = S_d$ is an initial segment of the cube order. Let y be the element in the cube order such that $y > \bar{x}$ and there is no other element between \bar{x} and y . Then $y_i = 0$ for $i \neq m_d + 1$, $y_{m_d+1} = k_d$. Clearly y is the lowest element in $P(C_{d-1}(n_{d-1}))$ in the cube order. Our assumption that $C_{d-1}(n_{d-1})$ is an initial segment of the cube order guarantees that $P(C_{d-1}(n_{d-1}))$ forms a complete

segment of the cube order starting from y . Therefore $C_d(n_d)$ is an initial segment of the cube order. By Proposition 2.1, we obtain the result. \square

THEOREM 3.3. *Let \mathcal{C} be a d -dimensional grid graph with n vertices and n_d be a positive integer such that n_d divides n . Then the total surface area of an equi-partition of \mathcal{C} into subgraphs of cardinality n_d is at least*

$$\frac{2n}{n_d} \sum_{r=1}^d [(r - m_r)(k_r + 1)^{m_r} k_r^{r-m_r-1} + m_r(k_r + 1)^{m_r-1} k_r^{r-m_r}]$$

where k_i and m_i , $i = 1, 2, \dots, d$ are calculated as mentioned in the definition of the quasi cube.

Proof. Since the equi-partition has n/n_d grid graphs and each graph is having n_d vertices, the result follows from Lemma 3.1 and Lemma 3.2. \square

3.1. Lower Bound Algorithm. Using the above results, this algorithm calculates a lower bound for the total surface area of an equi-partition of a grid graph with cardinality n into subgraphs of cardinality n_d .

```

begin { Lower Bound }
  for  $r = 1$  to  $d$ 
     $area \leftarrow 0$ 
     $k_r \leftarrow 0$ 
     $m_r \leftarrow 0$ 
  end for
   $s \leftarrow n_d$ 
   $r \leftarrow d$ 
  while ( $s > 0$ )
    Select  $k_r$  so that  $k_r^r \leq s < (k_r + 1)^r$ 
    Select  $m_r$  so that  $(k_r + 1)^{m_r} k_r^{r-m_r} < s \leq (k_r + 1)^{m_r+1} k_r^{r-m_r-1}$ 
     $s \leftarrow s - (k_r + 1)^{m_r} k_r^{r-m_r}$ 
     $area \leftarrow area + (r - m_r)(k_r + 1)^{m_r} k_r^{r-m_r-1} + m_r(k_r + 1)^{m_r-1} k_r^{r-m_r}$ 
     $r \leftarrow r - 1$ 
  end while
  return  $\frac{2n}{n_d} area$ 
end {Lower Bound }

```

When $n = n_d$, Lower Bound Algorithm returns the minimum surface area of a grid graph with cardinality n . The grid graphs that meet the minimum surface area are of special interest in many fields as mentioned in Section 1. Table 3.1 shows the minimum surface area of a grid graph for some selected values of n and d .

4. Equi-Partitioning Algorithm. Let K_1, K_2, \dots, K_d, V be positive integers and V divides $K_1 K_2 \dots K_d$. Here we present a heuristic algorithm for $MSA([K_1] \times [K_2] \times \dots \times [K_d], V)$ problem. The key idea of the algorithm is to decompose the given hyper-rectangle into a collection of well chosen “towers” of the form $[k_1] \times [k_2] \times \dots \times [k_{d-1}] \times [K_d]$, in which the tower “bases” $[k_1] \times [k_2] \times \dots \times [k_{d-1}]$ tile the $[K_1] \times [K_2] \times \dots \times [K_{d-1}]$ base of the given hyper-rectangle. Associated with each tower is the area of a particular equi-partition of the tower. In theory, one could construct an “optimal” tiling of the given base by minimizing the total of the surface areas of the corresponding towers, but this problem is comparable in

TABLE 3.1
The minimum surface area of a d -dimensional grid graph with n vertices

n	$d = 2$	$d = 3$	$d = 4$	$d = 5$	$d = 6$	$d = 7$	$d = 8$
1	4	6	8	10	12	14	16
10	14	30	50	70	90	110	130
10^2	40	130	262	416	576	768	968
10^3	128	600	1450	2538	3878	5392	7016
10^4	400	2800	8000	16066	26000	38506	51750
10^5	1266	12974	45262	100000	176900	273928	382822
10^6	4000	60000	253806	634564	1200000	1963298	2878960
10^7	12650	278708	1423044	3991582	8209238	14000000	21495344
10^8	40000	1293058	8000000	25152000	55892396	101208628	160000000

difficulty to the original problem. Our heuristic builds up the base tiling via easily computed partitions of lower dimensions. The main task of the algorithm involves solving $1 + K_1 + K_1K_2 + \dots + K_1\dots K_{d-2}$, $d \geq 2$ knapsack problems that represent total areas associated with “tilings” by towers. For $d = 2$, there is only one knapsack problem and this case was considered by Martin [8]. The algorithm is implemented in $d + 1$ phases.

In Phase-0, we partition each hyper-rectangle, $[k_1] \times [k_2] \times \dots \times [k_{d-1}] \times [K_d]$ where $k_i \in [K_i]$ ($1 \leq i \leq d - 1$), into components of size V and calculate the total surface area for the partition. If $V < k_1k_2\dots k_{d-1}$ (in which case a component created by the algorithm would be “badly shaped” by having size 1 in dimension d) or V does not divide $k_1k_2\dots k_{d-1}K_d$, we will consider the component as an invalid component and assign ∞ as the surface area. Otherwise we calculate the area of the partition of the tower assuming the vertices of each component are assigned by the *lexicographical order* on Z^d which can be described as follows. Let $x = (x_1, x_2, \dots, x_d)$ and $y = (y_1, y_2, \dots, y_d)$ be two d -tuples in $[k_1] \times [k_2] \times \dots \times [k_{d-1}] \times [K_d]$. We say $x > y$ if $j = \max\{i : x_i - y_i \neq 0\}$ and $x_j - y_j > 0$. If $x > y$ then we assign y before x . If the component size is V , we may assign the first V cells in the above total order to the first component and the next V cells to the second component and carry on the process until we fill the targeted region.

For a “tall” tower whose base is appropriately chosen, lexicographic assignment generates components that are good approximations to quasi-cubes and hence near-optimal surface areas are obtained. This property is addressed in the following two theorems. Theorem 4.1 presents an expression that may be used to compute the total surface area resulting from lexicographic assignment of the nodes in a tower under the assumptions that the tower has an integral number of components and that each component has a “height” of at least one. Theorem 4.2 establishes under some mild assumptions that the average surface area of the components produced by lexicographic assignment of nodes in “tall” towers with “well-chosen” bases asymptotically approaches minimum surface area. The computational results presented in Section 5 demonstrate this asymptotic optimality property of lexicographic assignment.

THEOREM 4.1 (Total area for lexicographic assignment). *Assume that V divides $k_1k_2\dots k_{d-1}K_d$, $V \geq k_1k_2\dots k_{d-1}$, and let $p \geq 1$ be the number of components of cardinality V in the d -dimensional region $[k_1] \times [k_2] \times \dots \times [k_{d-1}] \times [K_d]$. Let $s_j = k_1k_2\dots k_j$ where $1 \leq j \leq d - 1$. Let r_{ij} be the j -dimensional remainder of the i th component which is calculated by $r_{ij} = iV \bmod s_j$, where $1 \leq i \leq p - 1$, and $1 \leq j \leq d - 1$. Then the total surface area of an equi-partition of $[k_1] \times [k_2] \times \dots \times [k_{d-1}] \times [K_d]$*

into p components of cardinality V using the lexicographical order is

$$(4.1) \quad 2[ps_{d-1} + \sum_{j=1}^{d-1} \frac{s_{d-1}}{k_j} K_d + \sum_{i=1}^{p-1} \sum_{j=1}^{d-1} a_{ij}],$$

where a_{ij} is calculated as follows.

$$a_{i1} = \begin{cases} 1 & \text{if } r_{i1} \neq 0, \\ 0 & \text{otherwise.} \end{cases}$$

When $j > 1$,

$$a_{ij} = \begin{cases} 0 & \text{if } k_j = 1, \\ r_{ij} & \text{if } k_j > 1 \text{ and } r_{ij} < s_{j-1}, \\ s_j - r_{ij} & \text{if } k_j > 1 \text{ and } s_j - r_{ij} < s_{j-1}, \\ s_{j-1} & \text{otherwise.} \end{cases}$$

Proof. It is clear that the outer surface area of the grid graph $[k_1] \times [k_2] \times \dots \times [k_{d-1}] \times [K_d]$ is $2[s_{d-1} + \sum_{j=1}^{d-1} \frac{s_{d-1}}{k_j} K_d]$. To find the area of the inner boundaries, we need to add the area between the i th component and the $(i+1)$ th component for each $1 \leq i \leq p-1$. Since $V \geq s_{d-1}$, there are $p-1$ inner boundaries and each boundary is shared by exactly 2 components. So the total area of the projections of the inner boundaries of the components in the d th direction is $2(p-1)s_{d-1}$. For $1 \leq j \leq d-1$, let a_{ij} be the common boundary area between i th component and the $i+1$ th component in the j th direction. The possible cases for a_{ij} are mentioned in the statement of Theorem 4.1. Here we explain two cases and leave the more straightforward cases for the reader's verification. We can visualize $[k_1] \times [k_2] \times \dots [k_j]$ as k_j cross sections of $[k_1] \times [k_2] \times \dots [k_{j-1}]$ in the j th direction. These cross sections are assigned to the i th component in increasing order of j . When $r_{ij} < s_{j-1}$, the i th component is assigned r_{ij} vertices from the first $[k_1] \times [k_2] \times \dots [k_{j-1}]$ cross section in the j th direction, and hence $a_{ij} = r_{ij}$. When $s_j - r_{ij} < s_{j-1}$, the i th component is assigned r_{ij} vertices from the last $[k_1] \times [k_2] \times \dots [k_{j-1}]$ cross section in the j th direction, and hence $a_{ij} = s_j - r_{ij}$. Summing up the contributions in each direction, we get the area between the i th component and the $(i+1)$ th component as $2 \sum_{j=1}^{d-1} a_{ij}$. So the total surface area of inner boundaries is equal to $2(p-1)s_{d-1} + 2 \sum_{i=1}^{p-1} \sum_{j=1}^{d-1} a_{ij}$. Adding the outer and inner boundary areas gives us our result. \square

THEOREM 4.2. *Consider the family of d -dimensional towers $T(N)$ of the form $[k_1(N)] \times [k_2(N)] \dots \times [k_{d-1}(N)] \times [K_d(N)]$, where $|k_i(N) - N| \leq C$ for some constant C for $1 \leq i \leq d-1$ and all $N = 1, 2, \dots$ and $K_d(N) = N^z$ where $z > 1$. Let $V = N^d$ and let $a_l(N^d)$ be the average surface area of the components of $T(N)$ generated lexicographically under the assumptions of Theorem 4.1. Then the average surface area $a_l(N^d)$ asymptotically approaches the minimum surface area (for volumes of size N^d), i.e.,*

$$\lim_{N \rightarrow \infty} \frac{a_l(N^d)}{a^*(N^d)} = 1$$

where $a^*(N^d)$ is the minimum surface area ($2dN^{d-1}$) for volume V .

Proof. By Theorem 4.1,

$$a_l(N^d) = 2s_{d-1}(N) + \frac{2}{p(N)} \left[\sum_{j=1}^{d-1} \frac{s_{d-1}(N)}{k_j(N)} K_d(N) + \sum_{i=1}^{p(N)-1} \sum_{j=1}^{d-1} a_{ij}(N) \right] \text{ where}$$

$s_{d-1}(N) = k_1(N)k_2(N)\dots k_{d-1}(N)$ and $p(N) = s_{d-1}(N) \times N^z/N^d$. Since $z > 1$, note that $p(N) \rightarrow \infty$ when $N \rightarrow \infty$. For sufficiently large N , it is easily verified that $a_l(N^d) \leq 2(N+C)^{d-1} + 2 \sum_{j=1}^{d-1} [\frac{N^d}{N-C} + d(N+C)^{d-2}]$. Thus $\limsup_{N \rightarrow \infty} \frac{a_l(N^d)}{a^*(N^d)} \leq 1$, but since $\frac{a_l(N^d)}{a^*(N^d)} \geq 1$ for all N , the conclusion follows. \square

4.1. Phase-0 (Surface area computation for towers). Surface area computations using formula (4.1) are performed for all "valid" towers.

begin {Phase-0 }

Define a $(d-1)$ -dimensional array $A_{d-1}[K_1][K_2]\dots[K_{d-1}]$ of surface areas as follows:

$V \leftarrow$ the cardinality of a component.

for $k_1 = 1$ to K_1

for $k_2 = 1$ to K_2

.

.

.

for $k_{d-1} = 1$ to K_{d-1}

begin

if V does not divide $k_1k_2\dots k_{d-1}K_d$ or $V < k_1k_2\dots k_{d-1}$

$A_{d-1}[k_1][k_2]\dots[k_{d-1}] \leftarrow$ INFINITY

else

$A_{d-1}[k_1][k_2]\dots[k_{d-1}] \leftarrow$ The total surface area given by (4.1) for the partition of $k_1 \times k_2 \times \dots \times k_{d-1}K_d$ into components of cardinality V using the lexicographical order

end if

end for

end {Phase-0 }.

4.2. Phase- i ($1 \leq i \leq d-1$) (Determine optimal base decompositions in successively higher dimensions). In the i th ($1 \leq i \leq d-1$) phase, we solve $K_0K_1K_2\dots K_{d-i-1}$ (where $K_0 = 1$) knapsack problems using dynamic programming. For each $k_i \in [K_i]$, We define the $k_1k_2\dots k_{d-i-1}$ th knapsack problem as follows.

minimize $\sum_{j=1}^{K_{d-i}} c_j x_j$

subject to $\sum_{j=1}^{K_{d-i}} h_j x_j = K_{d-i}$

where each x_j is a nonnegative integer variable, $h_j = j$, and

$c_j = A_{d-i}[k_1][k_2]\dots[k_{d-i-1}][j]$ (where A_{d-i} is a $(d-i)$ -dimensional matrix (i.e., $i = 1$ case is used in Phase-0)) contains the optimal objective value of the $k_1k_2\dots k_{d-i-1}j$ th knapsack problem when $i > 1$, and the area of the equi-partition of subdomain $[k_1] \times [k_2] \times \dots \times [k_{d-i-1}] \times [j] \times [K_d]$ when $i=1$ (from Phase 0).

The above knapsack problem determines the combination of hyper-rectangular subdomain heights in direction $(d-i)$ that minimizes the total surface area as expressed by the objective function. The value of x_j represents how many sub domains of height j in the $(d-i)$ direction (i.e., $[k_1] \times [k_2] \times \dots \times [k_{d-i-1}] \times [j] \times [K_{d-i+1}] \dots \times [K_d]$) are in the optimal solution. The reader can observe that Phase- i knapsack problems use the results of Phase- $(i-1)$ knapsack problems as their area measure objective

coefficients c_j . Initially, Phase-0 area values are used as c_j values for Phase-1.

```

begin { Phase- $i$  ( $1 \leq i \leq d-1$ ) }
  if(i= d-1) (i.e. the final knapsack problem)
     $A_0 \leftarrow$  The solution of the final knapsack problem
    (i.e. the optimal objective value and a corresponding solution  $\{(h_j, x_j)|x_j \neq 0\}$ )
    call Phase- $d$ 
    return
  else
    Define a  $d-i-1$  dimensional array  $A_{d-i-1}[K_1][K_2] \dots [K_{d-i-1}]$ .
    for  $k_1 = 1$  to  $K_1$ 
    for  $k_2 = 1$  to  $K_2$ 
    .
    .
    .
    for  $k_{d-i-1} = 1$  to  $K_{d-i-1}$ 
    begin
       $A_{d-i-1}[k_1][k_2] \dots [k_{d-i-1}] \leftarrow$  The solution of the  $(k_1 k_2 \dots k_{d-i-1})$ th
      Knapsack problem (i.e. the optimal objective value and a corresponding solution
       $\{(h_j, x_j)|x_j \neq 0\}$ )
    end for
  end if
end { Phase- $i$  }

```

Our equi-partitioning algorithm is feasible when the above Phase- $d-1$ algorithm which solves the last knapsack problem is feasible. The following theorem guarantees the feasibility of Phase- $(d-1)$ knapsack problem.

THEOREM 4.3. *If V divides $K_1 K_2 \dots K_d$ then Phase- $(d-1)$ knapsack problem is feasible.*

Proof. Since V divides $K_1 K_2 \dots K_d$, we can find positive integers p_i ($1 \leq i \leq d$) such that p_i divides K_i and $V = p_1 p_2 \dots p_d$ (i.e., use the fundamental theorem of algebra). Now we can write $[K_i] = \cup_{j=0}^{n_i} j p_i + [p_i]$, where $n_i = (K_i/p_i) - 1$. We consider the $p_1 p_2 \dots p_{d-i-1}$ th knapsack problem in Phase i ($i < d-1$) where $c_j = A_{d-i}[p_1][p_2] \dots [p_{d-i-1}][j]$. This problem has a feasible solution $x_{p_{d-i}} = n_{d-i}$, and $x_j = 0$ if $j \neq p_{d-i}$ with the objective value $c_{p_{d-i}} n_{d-i}$ if $c_{p_{d-i}}$ is finite. Now we show that $c_{p_{d-i}} = A_{d-i}[p_1][p_2] \dots [p_{d-i-1}][p_{d-i}]$ is finite for each i ($1 \leq i \leq d-1$) by induction on i . In our base case ($i = 1$), since V divides $p_1 p_2 \dots p_{d-1} K_d$ and $p_1 p_2 \dots p_{d-1} \leq V$, we get a finite value for $c_{p_{d-1}} = A_{d-1}[p_1][p_2] \dots [p_{d-1}]$ (from Phase 0) which is the total surface area of the partition of the hyper-rectangle $[p_1] \times [p_2] \times \dots [p_{d-1}] \times K_d$. We assume $c_{p_{d-i}}$ is finite for ($i < d-1$). Then $p_1 p_2 \dots p_{d-i-1}$ th knapsack problem in Phase- i ($i < d-1$) is feasible as mentioned before and $c_{p_{d-i-1}}$ gets its finite optimal objective value. Therefore, by induction, $c_{p_{d-i}}$ is finite for all i ($1 \leq i \leq d-1$). Since c_{p_1} is finite, Phase- $(d-1)$ knapsack problem is feasible with a feasible solution of $x_{p_1} = n_1$, and $x_j = 0$ if $j \neq p_1$ with the objective value $c_{p_1} n_1$. \square

4.3. Phase- d (Assignment). Let $x = (x_1, x_2, \dots, x_{n_j})$ and $h = (h_1, h_2, \dots, h_{n_j})$ represent the solution of a knapsack problem that partitions the j th direction such that $\sum_{i=1}^{n_j} h_i x_i = K_j$ and each $x_i > 0$. Now we can partition $[K_j]$ such that $[K_j] = \cup_{i=1}^{m_j} k_{ji} + [H_{ji}]$ where $m_j = \sum_{i=1}^{n_j} x_i$, $k_{ji} = \sum_{p=1}^{q-1} x_p h_p + (i-1)h_q$ and $H_{ji} = h_q$

if $1 + \sum_{p=1}^{q-1} x_i \leq i \leq \sum_{p=1}^q x_i$.

```

begin {Phase-d }
  count ← 0
  proc ← 1
  for  $i_d = 0$  to  $K_d - 1$  begin Select the solution  $(x, h)$  stored in  $A_0$  from Phase-( $d-1$ )
  which partitions  $[K_1]$ 
  for  $i_1 = 1$  to  $m_1$ 
  begin
  Select the solution  $(x, h)$  stored in  $A_1[H_{1i_1}]$  from Phase-( $d-2$ ) which
  partitions  $[K_2]$ 
  for  $i_2 = 1$  to  $m_2$ 
  begin
  .
  .
  .
  Select the solution  $(x, h)$  stored in  $A_{d-2}[H_{1i_1}][H_{2i_2}] \dots [H_{(d-2)i_{(d-2)}}]$  from
  Phase-1 which partitions  $[K_{d-1}]$ 
  for  $i_{d-1} = 1$  to  $m_{d-1}$ 
  begin {assignment }
  for  $j_d = 0$  to  $K_d - 1$ 
  for  $j_{d-1} = k_{(d-1)i_{(d-1)}} to  $k_{(d-1)i_{(d-1)}} + H_{(d-1)i_{(d-1)}} - 1$ 
  .
  .
  .
  for  $j_1 = k_{1i_1}$  to  $k_{1i_1} + H_{1i_1} - 1$ 
  grid[ $j_1$ ][ $j_2$ ] ... [ $j_d$ ] ← proc
  count = count + 1
  if(count =  $V$ ) then
  proc ← proc + 1
  count ← 0
  end if
  end for (all  $j$ )
  end {assignment}
  end for (all  $i$ )
end {Phase-d}$ 
```

4.4. Analysis of the Equi-partitioning Algorithm. First we show that our algorithm solves the $MSA([K_1] \times [K_2] \times \dots [K_d], V)$ problem in $O(n)$ ($n = K_1 K_2 \dots K_d$) time if each K_i is in $O(n^{1/d})$. In Phase 0, we do n/K_d assignments and each assigned value takes at most $O(n^{1/d})$ time to calculate (Theorem 4.1). Since $K_d = O(n^{1/d})$, we need $O(n)$ time for Phase 0. In Phase- i ($0 \leq i \leq d-1$), we solve $K_0 K_1 \dots K_{d-i-1}$, ($K_0 = 1$), knapsack problems using dynamic programming which takes $O(K_{d-i}^2) = O(n^{2/d})$ time for each problem. Hence we solve Phase- i in $O(n^{(d-i+1)/d})$ time with the worst case $i = 1$ giving $O(n)$. It is clear that the final assignment phase (Phase- d) also takes $O(n)$ time. Therefore the algorithm runs in $O(n)$ time when each K_i is $O(n^{1/d})$.

Let us define a *gap* of a feasible solution for the MSA problem as the difference between a lower bound and an upper bound for the total area of the partition. The




FIG. 5.1. $Gap\%$ versus the number of vertices (M^3) for $MSA([M]^3, M)$ ($1 \leq M \leq 1000$)




FIG. 5.2. Execution time versus the number of vertices (M^3) for $MSA([M]^3, M)$ ($1 \leq M \leq 1000$)

percent gap ($gap\%$) is the gap as a percentage of the given lower bound. Although the above algorithm is always feasible when V divides $K_1 K_2 \dots K_d$, there are examples where the algorithm may not perform well (i.e., large $gap\%$). For example, consider $MSA([P]^4, P^2)$ problem where P is a prime. In this problem every valid domain in Phase-0, $[k_1] \times [k_2] \times [k_3] \times [P]$, has exactly one of k_1, k_2 or k_3 equal to P . So the area of each component is $\Omega(P^{4/3})$. The lower bound given in Theorem 3.2 for the area of a component is $O(P)$. Hence $gap\%$ is $\Omega(P^{1/3})$ which is undesirable for large P . We can modify Phase-0 and Phase- d of our algorithm to avoid these undesirable gaps. In Phase-0, when V does not divide $k_1 k_2 \dots k_{d-1} K_d$ (i.e., partition contains a partial component), the approach of Theorem 4.1 can be used to obtain an upper bound for the area of the partition. Instead of assigning ∞ to $A_{d-1}[k_1][k_2] \dots [k_{d-1}]$, we assign $U + 2s_{d-1}$ where U is the total area expression in Theorem 4.1 assuming $p = \lceil k_1 k_2 \dots k_{d-1} K_d / V \rceil$. In the final phase, Phase- d , we first assign labels for the full components and then work with the remaining partial components. We get at most one inner boundary inside a partial component justifying our upper bound $U + 2s_{d-1}$ in Phase-0.

5. Computational Results. The algorithms were coded in C and ran in 3.0GHz PCs with Linux OS.

We present computational results for selected three and four dimensional problems. Figure 5.1 shows the results for $MSA([M]^3, M)$ ($1 \leq M \leq 1000$). The percent-




FIG. 5.3. *Gap%* versus the number of vertices (M^4) for $MSA([M]^4, M^2)$ ($1 \leq M \leq 400$)




FIG. 5.4. *Execution time* versus the number of vertices (M^3) for $MSA([M]^4, M^2)$ ($1 \leq M \leq 400$)

age of solutions that are at the lower bound (i.e., optimal solutions) is 10.3%, and within 4% of gap is 95.3%. Figure 5.2 shows the execution times for the cases given in Figure 5.1. Due to memory limitation for larger problems, we have excluded the final assignment phase. So the running times in the graph are for obtaining the solutions without actual assignments.

Figure 5.3 shows the results for $MSA([M]^4, M^2)$ ($1 \leq M \leq 400$). The percentage of solutions that are at the lower bound (i.e., optimal solutions) is 15.0%, and within 4% of gap is 77.75%. Here we used the modified algorithm as mentioned in Section 4.4. Figure 5.4 shows the execution times for the cases given in Figure 5.3.

6. Conclusions and Future Research. We have developed a heuristic that runs in linear time in the size of the graph (for well shaped hyper-rectangles) that generates asymptotically optimal equi-partitions of hyper-rectangles, and demonstrated this performance for graphs of size up to 10^{10} .

When the area of an equi-partition does not meet the lower bound, it is very challenging to decide whether the partition is optimal. As an example, one may consider the equi-partition for the $MSA([5]^3, 5)$ problem shown in Figure 1.1 which is the smallest non trivial problem in this family. This partition has a surface area of 502, bound gap of 2, and with only one non-optimal component which is shown with the letter 'A'. Is this an optimal solution? Constraint programming may allow

optimality verification for smaller problems such as this.

Other interesting extensions to this research include relaxing the divisibility assumptions (i.e., modified algorithm), and improving the assignments obtained by more complex procedures for bad components as suggested in [5] for the two-dimensional case.

Acknowledgments. The authors thank Winston Yang for bringing the references [1] and [3] to their attention.

REFERENCES

- [1] L. ALANZO, AND R. CERF, *The three dimensional polyominoes of minimal area*, Electronic Journal of Combinatorics, 3.1 (1996), R27 pp. 1-39.
- [2] S. L. BEZRUKOV, AND B. ROVAN, *On partitioning grids into equal parts*, Computers and Artif. Intell., 16 (1997), pp. 153–165.
- [3] B. BOLLABAS, AND M. LEADER, *Edge Isometric Inequalities in the Grid*, Combinatorica, 11.4 (1991), pp. 299–314.
- [4] I. T. CHRISTOU, AND R. R. MEYER, *Optimal equi-partition of rectangular domains for parallel computation*, Journal of Global Optimization, 8 (1996), pp. 15–34.
- [5] W. W. DONALDSON, *Grid-graph partitioning*, Ph.D. thesis, Department of Computer Science, University of Wisconsin-Madison, Madison, WI, 2000.
- [6] M. R. GAREY AND D. S. JOHNSON, *Computers and Intractability: A Guide to the Theory of NP-Completeness*, W. H. Freeman, 1979.
- [7] S. GHANDEHARIZADEH, R. R. MEYER, G. SCHULTZ, AND J. YACKEL, *Optimal balanced partitions and a parallel database application*, ORSA Journal on Computing, 4 (1993), pp. 151–167.
- [8] W. MARTIN, *Fast equi-partitioning of rectangular domains using stripe decomposition*, Discrete Appl. Math., 82 (1998), pp. 193–207.
- [9] E. J. NEVES, *A discrete variational problem related to ising droplets at low temperatures*, Journal of Statistical Physics, 80 (1995), pp. 103–123.
- [10] R. J. SCHALKOFF, *Digital Image Processing and Computer Vision*, John Wiley & Sons, Inc., 1989.
- [11] J. STRIKWERDA, *Finite Difference Schemes and Partial Difference Equations*, Wadsworth & Brooks, 1989.
- [12] J. YACKEL, *Minimum-perimeter tiling for optimization of parallel computation*, Ph. D. thesis, Department of Computer Science, University of Wisconsin-Madison, Madison, WI, 1993.
- [13] J. YACKEL, *Partitioning grid graphs with partial cubes*, private communication.
- [14] J. YACKEL, R. R. MEYER, AND I. CHRISTOU, *Minimum perimeter domain assignment*, Mathematical Programming, 78 (1997), pp. 283–303.